

FIFRA §2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA, ARIZONA, ARKANSAS, CALIFORNIA, COLORADO, CONNECTICUT, DELAWARE, DISTRICT OF COLUMBIA, FLORIDA, GEORGIA, HAWAII, IDAHO, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, LOUISIANA, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA, MISSISSIPPI, MISSOURI, NEBRASKA, NEVADA, NEW HAMPSHIRE, NEW JERSEY, NEW MEXICO, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, UTAH, VERMONT VIRGINIA, WASHINGTON, WEST VIRGINIA, AND WISCONSIN



## EPA REG. NO: 10163-357

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- This labeling contains supplemental instructions that may not appear on the product label. All applicable directions, restrictions, and precautions on the product label are to be followed.
- This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the United States Environmental Protection Agency (EPA).

### PRODUCT INFORMATION

EcoSwing is a biofungicide for use on growing crops in agricultural, greenhouse, ornamental and turf use sites to control fungal diseases such as powdery mildew (*Oidium spp.*), rust, and *Botrytis cinerea*. Optimum disease control is achieved when the biofungicide is used according to label directions and applied in a regularly scheduled preventative spray program.

The use of EcoSwing is compatible with the principles of Integrated Pest Management (IPM) and programs that attempt to minimize disease resistance to fungicides. Unlike single-site mode of action fungicides which are at risk from disease resistance, EcoSwing has a multi-site mode of action, and may be used to delay or prevent the development of resistance to single site fungicides. Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of EcoSwing in IPM and resistance management programs.

**PRE-HARVEST INTERVAL:** EcoSwing has a zero day pre-harvest interval and can be applied up to and including the day of harvest

### APPLICATION DIRECTIONS

CROP	PEST	COMMENTS
<b>Ornamental Plants, Shrubs, and Trees (including those grown in landscapes)</b> Such as: Amaranthus, Aster, Azalea, Caladium, Carnation, Chrysanthemum, Dahlia, Daisy, Ferns, Ficus, Fuschia, Gardenia, Impatiens, Iris, Ivy, Jasmine, Lilac, Lilies, Marigold, Philodendron, Poinsettia, Rose, Zinnia	Powdery Mildew, Rust, Botrytis	For fogging applications, apply 1.5 – 2 pts. per acre or 0.55-0.73 fl oz per 1000 sq. feet.  Applications can be made in green houses, with automatic fogger equipment. Apply the same amount of EcoSwing® per unit area (acre or 1000 sq ft) as would be applied in a dilute spray volume and adjust water volume per the specifications of the chemical fogger manufacturer.  <b>DO NOT</b> apply through fogger equipment when workers are present in the greenhouse during the application.

EcoSwing® is a registered trademark used under license by Gowan Company, L.L.C.

Produced For:  
 Gowan Company, LLC  
 PO Box 5569  
 Yuma, AZ 85366-5569

