



FIFRA Section 24(c) Special Local Need (SLN) Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

Fulfill® Insecticide

For Control of Aphids on Listed Herb and Vegetable Crops Grown for Seed

EPA Reg. No. 66222-274

EPA SLN No. WA-190003

This label for Fulfill Insecticide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2024.

Directions for Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PEST CONTROL, OR ILLEGAL RESIDUES.

Crops - Herb and vegetable crops grown for seed: Arugula; Beet, garden; Beet, sugar; Broccoli, Chinese; Broccoli raab; Brussels sprouts; Cabbage; Carrot; Cauliflower; Cabbage, Chinese, bok choy; Cabbage, Chinese, napa; Collards; Endive; Kale; Kohlrabi; Lettuce, head; Lettuce, leaf; Mustard greens; Mustard, seed; Parsley; Parsnip; Radish (other than radish, Oriental); Rape greens; Rutabaga; Spinach; Spinach mustard; Swiss chard; Turnip.

Pests:

Bean aphid, Black bean aphid, Cabbage aphid, Green peach aphid, Lettuce aphid, Melon aphid, Pea aphid, Potato aphid, Turnip aphid, Willow-carrot aphid.

Fulfill Insecticide works primarily by ingestion, but also has some contact activity. Affected aphids stop feeding shortly after exposure but may remain on the plant foliage until they die, which is usually within 4-10 days.

Application Rate:

Foliar application: 2.75 oz/A.

Application Concentration / Methods / Frequency / Timing:

Use a minimum of 5 gals of water/A when applied by air and a minimum of 10 gals of water/A when applied by ground.

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage. The addition of a penetrating type spray adjuvant, at the manufacturer's suggested rate, is recommended to provide optimum coverage and penetration. Apply when aphids first appear, before populations build to damaging levels. Two applications may be needed to control persistent aphid populations.

Restrictions/Precautions:

- Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- Do not apply more than 2.75 oz/A per application.
- Do not exceed a total of 5.5 oz/A per crop per season.
- Allow a minimum of 7 days between applications
- Allow a minimum of 14 days (PHI) between the last application and harvest.
- For use under this SLN label do not apply this product through any type of irrigation system.
- Do not apply Fulfill Insecticide directly to bees that are actively foraging in the field.

Crops Grown for Seed:

1. All herb and vegetable seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
2. No portion of the herb and vegetable seed plants, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
3. All seed from herb and vegetable seed crops shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
4. No seed from herb and vegetable seed crops may be distributed for human consumption or animal feed.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at: <https://agr.wa.gov/departments/pesticides-and-fertilizers/pesticides/waste-pesticide>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington

Fulfill is a registered trademark of ADAMA
05292019

Section 24(c) Registrant:
Makhteshim Agan of North America, Inc. d/b/a ADAMA
3120 Highwoods Blvd., Suite 100
Raleigh, NC 27604

